


We have taken great care in collecting and checking the information provided in this leaflet to make sure that it is accurate.

We do not take responsibility for any inaccurate or missing information in this leaflet. However, if you have any concerns about this information, please contact us.



Cancer Screening Programmes

Birmingham Women's
NHS Foundation Trust



Ref 158/07/2011
Review 07/2014



© West Midlands
Cervical Screening QA
Reference Centre
Last updated June 2007

Birmingham Women's 
NHS Foundation Trust

Colposcopy Department, 3rd Floor, Metchley Park Road
Edgbaston, Birmingham B15 2TG
Colposcopy Reception Tel: 0121 607 4733
Fax: 0121 607 4706 Website: www.bwhct.nhs.uk



More information

If you have any more questions about your referral or treatment, phone our clinic on **0121 472 1377** ext 2698 or 4705

Contact name:

Debbie Wise (Clinical Nurse Specialist)
Sam Venner (Clinical Nurse Specialist)

You may also contact your family doctor and they will be happy to help you.

You can find more information on cervical screening on the National Cancer Screening website at www.cancerscreening.nhs.uk/cervical/index.html

2 Your cervical screening test has found evidence of possible abnormal cells.

You have been asked to come for a further examination because your cervical screening test has found abnormal changes in the cells of the cervix (the neck of the womb). These abnormal changes are known as **dyskaryosis**. This is not unusual, about one in 20 cervical screening tests are reported as abnormal. Please try not to worry. The NHS Cervical Screening Programme aims to pick up these changes so that we can take action at an early stage.

What is dyskaryosis? This is the term given to the abnormal changes that can act as an early warning that cervical cancer might develop in the future. It is important to remember that it is **very rare** indeed for these abnormal cells to be cancer.

For many women, their abnormal result will show **borderline changes or mild dyskaryosis**. These are small, low-grade changes in the cells and will very often return to normal by themselves. It is often safe to keep an eye on the situation to see if this happens rather than having immediate treatment.

For some women, their result will show **moderate or severe dyskaryosis**. These changes are not cancer, and in most cases do not lead to cancer in the future. However, these changes are less likely to return to normal by themselves and usually act as an early warning sign that, over time, cervical cancer may develop. Because of this, they will usually need simple outpatient treatment that is virtually 100% effective in most cases.

3 Other signs or symptoms not related to having a cervical screening test.

You may not have been referred for a colposcopy because of a cervical screening test. However, you may have been experiencing signs or symptoms which may be related to the health of your cervix. The colposcopy clinic is the best place to have these symptoms checked.

Is there anything that I need to know before I have a colposcopy?

- There is no need to avoid having sex before you have a colposcopy examination, unless you have a coil fitted.
- You can eat and drink as normal before your appointment.
- Bring a list of any medication you are taking and let the colposcopist know of any medical history or allergies.
- Please also make a note of when you had your last period.
- You will be asked to remove clothes below the waist. Because of this, you may find it more comfortable to wear a full skirt or dress, which you would not need to remove for the examination.
- If you need treatment during your appointment, you may need to remove any jewellery you wear below the waist (such as belly button or vaginal piercings). It is important that you are aware of this so you can remove the jewellery before you come for your appointment.
- After the examination, you may have some discharge from your vagina. Although we will provide pads, you are more than welcome to bring panty liners with you if you prefer.
- You should be able to drive home afterwards but for car-insurance purposes, you should ask your colposcopist whether you are fit to drive after your treatment. It may be wise to bring someone with you in case you do not feel well after your appointment.

Practical questions

What should I do if I have a period on the day of my colposcopy?

Ring the clinic and check that you should go ahead with your appointment, as this often depends on what your appointment is for. If you feel awkward about having an examination during your period, contact the clinic and change the time of your appointment.

Can I have sex in the week before I have a colposcopy?

You do not need to avoid having sex before you have a colposcopy, unless you have a coil fitted. If this is the case, please use an extra method of contraception for seven days before your appointment, as sometimes it is necessary to remove the coil before you have the treatment.

What happens if I am pregnant or think I might be?

It is important that you keep your colposcopy appointment. A colposcopy can be carried out quite safely during pregnancy, but any investigations will usually be postponed until after the baby is born. When you come for your appointment, we can talk to you more about this.

Will having a colposcopy affect my fertility or sex life?

Having a colposcopy does not affect either of these. However, different treatments may have different effects and we will be happy to discuss this with you at your appointment. If you have any treatment, you will have to avoid having sex for four weeks afterwards. If you have a special event coming up (for example, you are getting married or going on holiday very soon after your appointment), please contact the clinic and we will discuss whether it is necessary to rearrange your appointment.

- It is normal to have some bloodstained discharge after treatment. If you have bloodstained discharge from the treatment, do **not** use tampons. Instead, use pads or panty liners. You should not use tampons until four weeks after the treatment. Do not worry if you do not have any discharge.
- You should avoid heavy lifting or vigorous exercise for a day or so after treatment. You should also avoid swimming for at least two weeks or until the discharge has stopped because of the risk of infection.
- If you have any questions after your treatment, please contact the colposcopy clinic on 0121 4721377 extension 2698 or 4705.
- We will send you a letter with your results. If you have not received this letter eight weeks after your appointment, please contact the clinic.

Rare problems with treatment

A small number of patients may suffer a pain in their pelvis or heavy bleeding from the treated area. This may happen immediately after treatment or within the next couple of weeks and may mean you have an infection, particularly if you have a temperature or strong-smelling discharge. It is very important that you contact your family doctor or the colposcopy clinic as you may need another examination or antibiotics.

Will I need to have check-ups?

Yes. It is important to keep your appointments to make sure that your cervix is healthy. Most colposcopy clinics recommend that you have a follow-up check between six and 12 months after the examination or treatment, although this will depend on the results of your examination. The check-ups may be back at the colposcopy clinic or could be at your family doctor's surgery or local clinic. We will discuss the exact details with you. **It is very important to complete the follow-up programme as this allows us to check that your cervix stays healthy.**

What will happen during my appointment?

When you arrive at the clinic, we will aim to see you as close to your appointment time as possible, but please be patient if there are delays that we could not have expected beforehand. You will then be called for your colposcopy, where you will meet the colposcopist (who may be a doctor or nurse) and the nurses who will be with you during the examination.

During your colposcopy appointment, the doctor or nurse will take down your medical history and explain the procedure to you. Following this, the colposcopist will carry out the examination. How long your appointment will take depends on whether any procedures are necessary but it is usually no longer than 20 minutes to half an hour.

What is a colposcopy?

Having a colposcopy is similar to having a cervical screening test. You will lie on a special couch that has supports for your legs. The colposcopist will warm and lubricate a speculum (a special instrument) and place it in your vagina (the same as when you had a cervical screening test). This will allow them to see your cervix. The colposcope (which, put simply, is binoculars on a stand) gives the colposcopist an enlarged 3-D view of the cervix. **The colposcope does not enter the vagina.**

The colposcopist will put a liquid onto the cervix with a cotton-wool ball – this may be cold and sometimes sting a little. This liquid highlights any abnormal cells and allows the colposcopist to make a diagnosis. If everything is normal, you may be allowed to go home straight away. However, the diagnosis may mean that it is necessary to take a **punch biopsy** or carry out **treatment**.

What is a punch biopsy?

A **punch biopsy** is a tiny piece of tissue taken from the cervix and sent to a pathologist so that it can be examined closely to see if there are changes in the cells. You will not normally need a local anaesthetic. The biopsy is about the size of a grain of rice. You may have a brown or red discharge from your vagina for the next few days after this. You may also feel discomfort like period pain for the next few hours. You should take your normal painkillers to ease this pain, but do not take more than recommended dose.

What if I need treatment?

The aim of **treatment** is to destroy or remove abnormal cells and allow healthy cells to replace them. Most types of treatment can be done as an outpatient in the colposcopy clinic on your first appointment, so you may want to arrange a fairly quiet day in case you are offered this. If you have a coil fitted, please use an extra form of contraception for seven days before your appointment, as sometimes it is necessary to remove the coil before you have the treatment. The treatment at this clinic is usually loop diathermy or cryocautery.

Types of treatment

Loop diathermy

This treatment uses an electric current to remove abnormal tissue from your cervix. A local anaesthetic (similar to that you would have at the dentist) is used to numb the cervix, and the abnormal cells are then lifted off the cervix using a wire loop. The colposcopist may then apply mustard-coloured paste to seal the area. During the procedure, you will hear various noises from the machinery. The examination and treatment lasts roughly 15 minutes. Afterwards, you will be given a sanitary pad (or you may want to bring your own) and asked to rest for a short time before leaving.

Cryocautery

Cryocautery is a procedure which destroys the tissue on the cervix by freezing it with nitrous oxide. It is usually performed to reduce the symptoms of excessive discharge or bleeding after having sex but can also be used to treat low-grade changes to the cervix. The colposcopist will place a small probe on the cervix for two minutes, freezing the abnormal tissue. You should not feel any discomfort during the procedure. You may experience a watery discharge for two to four weeks after the treatment as the cervix heals.

What will the treatment show?

The technical term used to refer to cell changes confirmed by a biopsy or treatment is **cervical intra-epithelial neoplasia**, more commonly known as CIN.

To make distinctions between the various states of changes in the cells, doctors have developed an increasing scale from one to three according to how many of the cells are affected. So, treatment results will most commonly be CIN 1, CIN 2 or CIN 3. The results of this treatment will show what follow-up you will need.

What normally happens after treatment?

Most women will feel okay after treatment. In a **very small** number of cases, the following may happen.

- Some women feel dizzy or faint when they get up following the treatment. If this happens to you, the nurse will ask you to lie down and will keep you under close observation until you have fully recovered.
- If you have a punch biopsy, you may experience discomfort like period pain for the next few hours. You should take your normal painkillers to ease the pain but not take more than the recommended dose.