

## Are there any side effects to morphine PCA?

Yes, but they are usually minor.

It can often make you feel a bit sleepy so it's quite normal to have extra oxygen whilst you have the PCA. It can sometimes make you feel a bit itchy or 'spaced out'.

The main side effect is feeling or being sick. If this happens to you, please tell your nurse because we can treat this.

## Do I have to have PCA?

No. There are other ways of controlling your pain, for example, by injections into your leg. If you're unsure about having a PCA, the anaesthetist, ward nurse or acute pain sister will be happy to discuss it with you further.

## Who can I talk to about PCA?

Contact a Sister in Pre-Operative Assessment on 0121 627 2755 or, after admission, ask to speak to the Acute Pain Nurse.

Ref 33/06/2010 Review 06/2013

Author:

**Pam Snell** - Acute Pain Sister

Birmingham Women's   
NHS Foundation Trust

Edgbaston, Birmingham B15 2TG  
Tel: 0121 472 1377 Fax: 0121 627 2602

Birmingham Women's   
NHS Foundation Trust



# *Patient Controlled Analgesia or PCA*

**Patient Information**

## Introduction

We now understand that even after the same operation, no two women are alike and that a person's experience of pain is a very individual thing. PCA is one method of pain relief that takes this into account.

## What is PCA?

PCA stands for patient-controlled analgesia (analgesia is a medical term for pain relief) and is a way of giving you painkilling drugs.

It consists of a pump containing a strong painkiller (usually morphine) which is connected by a tube to the back of your hand or arm.

## If I decide to have PCA, what will I have to do?

When you wake up in recovery after your operation, you'll be given a small button to hold.

If you're in pain, all you have to do is press and release the button, you will hear a bleep and the PCA pump will give you a small dose of morphine.

When you get back to the ward, continue to press and release the button whenever you have any pain.

The day after your operation, you might find it useful to give yourself an extra dose or two before doing something that might make you feel uncomfortable, e.g. getting in and out of bed or before physiotherapy.

**It's important to know that it's only you who can press the button.**

## Could I give myself an overdose?

The pump has been programmed so that this cannot happen. You can press the button as many times as you like, but the pump will only give you a dose of morphine every 5 minutes and it will only let you have a certain amount every hour, so it is doubly safe.

## Could I become addicted to it?

No, you'll only need morphine for a day or two and as the pain gets better, you'll use the PCA less.

## Should I wait for the pain to build up before I press the button?

No. If your tummy hurts when you try to move, then that is a good sign that you should press the button. Everyone's pain is different, so press the button whenever you need to.

## Will the PCA take all my pain away?

No, you'll still experience some discomfort and mild pain.

## Does the PCA mean that I won't need any other painkillers?

No. We'll give you other painkillers, depending on what suits you, as well as the PCA. It's important you take these tablets regularly, especially after you've finished with the PCA.

## Does it mean that the nurses will pay less attention to me?

No. The nurses will regularly check that you are comfortable.

## Will the PCA stop me getting out of bed

No, the PCA pump is light and portable, so you can still sit out of bed.